


Class X
Unit: 8 (Trigonometry)
Revision Notes

Revision Notes on Introduction to Trigonometry

Trigonometry

To find the distances and heights we can use the mathematical techniques, which come under the **Trigonometry**. It shows the relationship between the sides and the angles of the triangle. Generally, it is used in the case of a right angle triangle.

Trigonometric Ratios



In a right angle triangle, the ratio of its side and the acute angles is the trigonometric ratios of the angles.

In this right angle triangle $\angle B = 90^\circ$. If we take $\angle A$ as acute angle then -

AB is the base, as the side adjacent to the acute angle.

BC is the perpendicular, as the side opposite to the acute angle.

AC is the hypotenuse, as the side opposite to the right angle.

Trigonometric ratios with respect to $\angle A$

Ratio	Formula	Short form	Value
$\sin A$	$\frac{\text{Perpendicular}}{\text{hypotenuse}}$	P/H	BC/AC
$\cos A$	$\frac{\text{Base}}{\text{hypotenuse}}$	B/H	AB/AC
$\tan A$	$\frac{\text{Perpendicular}}{\text{base}}$	P/B	BC/AB
$\operatorname{cosec} A$	$\frac{\text{Hypotenuse}}{\text{perpendicular}}$	H/P	AC/BC
$\sec A$	$\frac{\text{Hypotenuse}}{\text{base}}$	H/B	AC/AB
$\cot A$	$\frac{\text{Base}}{\text{perpendicular}}$	B/P	AB/BC

Remark

- If we take $\angle C$ as acute angle then BC will be base and AB will be perpendicular. Hypotenuse remains the same i.e. AC. So the ratios will be according to that only.
- If the angle is same then the value of the trigonometric ratios of the angles remain the same whether the length of the side increases or decreases.
- In a right angle triangle, the hypotenuse is the longest side so $\sin A$ or $\cos A$ will always be less than or equal to 1 and the value of $\sec A$ or $\operatorname{cosec} A$ will always be greater than or equal to 1.

Reciprocal Relation between Trigonometric Ratios

$\operatorname{Cosec} A$, $\sec A$, and $\cot A$ are the reciprocals of $\sin A$, $\cos A$, and $\tan A$ respectively.

$\sin \theta = \frac{1}{\operatorname{cosec} \theta}$	$\operatorname{cosec} \theta = \frac{1}{\sin \theta}$	$\sin \theta \operatorname{cosec} \theta = 1$
$\cos \theta = \frac{1}{\sec \theta}$	$\sec \theta = \frac{1}{\cos \theta}$	$\cos \theta \sec \theta = 1$
$\tan \theta = \frac{1}{\cot \theta}$	$\cot \theta = \frac{1}{\tan \theta}$	$\tan \theta \cot \theta = 1$

Quotient Relation

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

Trigonometric Ratios of Some Specific Angles


Trigonometry Table

	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined
$\operatorname{cosec} \theta$	Not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
$\sec \theta$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	Not defined
$\cot \theta$	Not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

Use of Trigonometric Ratios and Table in Solving Problems

Example

Find the lengths of the sides BC and AC in ΔABC , right-angled at B where $AB = 25$ cm and $\angle ACB = 30^\circ$, using trigonometric ratios.



Solution

To find the length of the side BC, we need to choose the ratio having BC and the given side AB. As we can see that BC is the side adjacent to angle C and AB is the side opposite to angle C, therefore

$$\tan C = \frac{AB}{BC}$$

$$\tan 30^\circ$$

$$\frac{25}{BC} = \frac{1}{\sqrt{3}}$$

$$BC = 25\sqrt{3} \text{ cm}$$

To find the length of the side AC, we consider

$$\sin 30^\circ = \frac{AB}{AC}$$

$$\frac{1}{2} = \frac{25}{AC}$$

$$AC = 50 \text{ cm}$$

Trigonometric Ratios of Complementary Angles

If the sum of two angles is 90° then, it is called **Complementary Angles**. In a right-angled triangle, one angle is 90° , so the sum of the other two angles is also 90° or they are complementary angles. so the trigonometric ratios of the complementary angles will be -

$$\sin (90^\circ - A) = \cos A,$$

$$\cos (90^\circ - A) = \sin A,$$

$$\tan (90^\circ - A) = \cot A,$$


$$\cot (90^\circ - A) = \tan A,$$

$$\sec (90^\circ - A) = \operatorname{cosec} A,$$

$$\operatorname{cosec} (90^\circ - A) = \sec A$$

Trigonometric Identities (Pythagoras Identity)

An equation is said to be a trigonometric identity if it contains trigonometric ratios of an angle and satisfies it for all values of the given trigonometric ratios.



In ΔPQR , right angled at Q, we can say that

$$PQ^2 + QR^2 = PR^2$$

Divide each term by PR^2 , we get

$$\frac{PQ^2}{PR^2} + \frac{QR^2}{PR^2} = \frac{PR^2}{PR^2}$$

$$\left\{\frac{PQ}{PR}\right\}^2 + \left\{\frac{QR}{PR}\right\}^2 = \left\{\frac{PR}{PR}\right\}^2$$

$$(\sin R)^2 + (\cos R)^2 = 1$$

$$\sin^2 R + \cos^2 R = 1$$

Likewise other trigonometric identities can also be proved. So the identities are-

$$\sin^2 R + \cos^2 R = 1$$

$$1 + \tan^2 R = \sec^2 R$$

$$\cot^2 R + 1 = \operatorname{cosec}^2 R$$

UNIQUESTUDYONLINE.COM